arXiv:1412.5837v1 [math.KT] 18 Dec 2014

Dennis trace map for certain K-groups of categories with cofibrations

Abhishek Banerjee

Hausdorff Institute for Mathematics, Poppelsdorfer Allee 45, D-53115, Bonn, Germany.
Email: abhishekbanerjee1313@gmail.com

Abstract

Let $\mathcal{C}$ be a small category with cofibrations. In this paper, we define the K-theory and Hochschild homology groups of $\mathcal{C}$ of order Y, where Y is an ordered finite simplicial set with basepoint. Further, we construct the Dennis trace map between these groups.

Résumé

Soit $\mathcal{C}$ une petite catégorie avec cofibrations. Dans cet article, nous définissons ses groupes de K-théorie et homologie de Hochschild d'ordre Y, où Y est un ensemble simplicial ordonné et fini avec point de base. De plus, nous construisons le morphisme de Dennis entre ces groupes.

1 Introduction

In [3], Waldhausen introduced categories with cofibrations and defined their K-theory groups by means of the "S-construction". For any $n \geq 0$, Waldhausen's S-construction associates to a small category $\mathcal{C}$ with cofibrations a category $S_n\mathcal{C}$. An object of $S_n\mathcal{C}$ is a chain of n composable cofibrations in $\mathcal{C}$ starting with the zero object (see (2.1)) Then, the objects of the simplicial category $S_\bullet\mathcal{C} = \{S_n\mathcal{C}\}_{n\geq 0}$ determine a simplicial set $S\mathcal{C} = \{obj(S_n\mathcal{C})\}_{n\geq 0}$. Then, the K-theory groups of the category $\mathcal{C}$ are defined to be the homotopy groups of the loop space $\Omega|S\mathcal{C}|$ of the geometric realization of the simplicial set $S\mathcal{C}$.

Let Ord_* denote the category of finite, totally ordered sets with basepoint. Let $SmCat_0$ denote the category of small categories with zero objects. The starting point for this article is the fact that the association $n \mapsto S_n\mathcal{C}$ extends naturally to a functor $S(\mathcal{C})$ from Ord_* to $SmCat_0$. Then, if we take a simplicial object $Y : \Delta^{op} \longrightarrow Ord_*$ of Ord_*, i.e., Y is an ordered finite simplicial set with basepoint, we consider the composition (see (2.4))

$$S^Y(\mathcal{C}) : \Delta^{op} \xrightarrow{Y} Ord_* \xrightarrow{S(\mathcal{C})} SmCat_0 \xrightarrow{obj} Sets_* \tag{1.1}$$

The purpose of this paper is to study the K-groups $K_p^Y(\mathcal{C})$, $p \geq 0$ of $\mathcal{C}$ of order Y which we define to be the homotopy groups of the loop space of the geometric realization of the simplicial set $S^Y(\mathcal{C})$ (see (2.5)). If Y, $Y' : \Delta^{op} \longrightarrow Ord_*$ are simplicially homotopy equivalent as simplicial objects of Ord_*, we show that $K_p^Y(\mathcal{C}) \cong K_p^{Y'}(\mathcal{C})$. Further, we describe a product structure $K_p^Y(\mathcal{C}) \times K_q^Y(\mathcal{D}) \longrightarrow K_{p+q}^Y(\mathcal{E})$ for a bi-exact functor $F : \mathcal{C} \times \mathcal{D} \longrightarrow \mathcal{E}$ (see (2.7)).

In the second part of the paper, we want to define Dennis trace maps from $K_p^Y(\mathcal{C})$ to appropriate Hochschild homology groups. For this, we consider the geometric realization $|CN(S^Y(\mathcal{C}))|$ of the bisimplicial set given by the cyclic nerve $CN(S^Y(\mathcal{C}))$ of the simplicial category $S(\mathcal{C}) \circ Y$. We define the Hochschild homology groups

$HH_p^Y(\mathcal{C})$ of $\mathcal{C}$ of order Y in terms of the singular homology of $|CN(S^Y(\mathcal{C}))|$ with coefficients in a given field k (see (3.3)). Again, if Y, $Y' : \Delta^{op} \longrightarrow Ord_*$ are simplicially homotopy equivalent as simplicial objects of Ord_*, we show that $HH_p^Y(\mathcal{C}) \cong HH_p^{Y'}(\mathcal{C})$. We show that the Hochschild homology groups also carry a product $HH_p^Y(\mathcal{C}) \otimes HH_q^Y(\mathcal{D}) \longrightarrow HH_{p+q+1}^Y(\mathcal{E})$ for a bi-exact functor $F : \mathcal{C} \times \mathcal{D} \longrightarrow \mathcal{E}$ (see (3.6)). Finally, for any $p \geq 0$, we construct a Dennis trace map $D_p^Y : K_p^Y(\mathcal{C}) \longrightarrow HH_p^Y(\mathcal{C})$.

2 The K-groups $K_p^Y(\mathcal{C})$ for a category with cofibrations

In this section and throughout this paper, we let $\mathcal{C}$ be a small category with cofibrations in the sense of Waldhausen [3]. In other words, $\mathcal{C}$ is a category with a zero object together with a subcategory $co\mathcal{C}$ satisfying the axioms (Cof1) and (Cof2) below. The morphisms in $co\mathcal{C}$ will be referred to as cofibrations and denoted by feathered arrows "$\rightarrowtail$".

(Cof1) Every isomorphism in $\mathcal{C}$ is a cofibration. For any object A in $\mathcal{C}$, the canonical morphism $0 \longrightarrow A$ is a cofibration.
(Cof2) Given a cofibration $A \rightarrowtail B$, its pushout $C \coprod_A B$ along any other morphism $A \longrightarrow C$ exists in $\mathcal{C}$ and the canonical morphism $C \longrightarrow C \coprod_A B$ is a cofibration.

Given a cofibration $A \rightarrowtail B$ in $\mathcal{C}$, its pushout $0 \coprod_A B$ along the morphism $A \longrightarrow 0$ will be denoted by B/A. The canonical morphism from B to the pushout $B/A = 0 \coprod_A B$ is referred to as a quotient map and denoted by $B \twoheadrightarrow B/A$. The sequence $A \rightarrowtail B \twoheadrightarrow B/A$ is referred to as a cofibration sequence. A functor between categories with cofibrations is said to be exact if it takes 0 to 0, preserves cofibrations as well as the pushout diagrams arising from axiom (Cof2).

Given $\mathcal{C}$ as above, we let $S_\bullet\mathcal{C}$ be the simplicial category associated to $\mathcal{C}$ by Waldhausen's S-construction (see [3, 1.3]). More explicitly, for any $n \geq 0$, an object of $S_n\mathcal{C}$ is a sequence $(a_0, a_1, ..., a_{n-1})$ of composable cofibrations:

$$0 = A_0 \overset{a_0}{\rightarrowtail} A_1 \overset{a_1}{\rightarrowtail} A_2 \overset{a_2}{\rightarrowtail} \dots \overset{a_{n-1}}{\rightarrowtail} A_n \tag{2.1}$$

together with a choice of quotients $A_{ij} = A_j/A_i$. Further, for any $i \leq j \leq k$, $A_{ij} \rightarrowtail A_{ik} \twoheadrightarrow A_{jk}$ is a cofibration sequence.

We now let Ord_* denote the category of finite totally ordered sets $(Z, *) = \{* < z_1 < z_2 < .... < z_k\}$ with basepoint $*$. A morphism $\phi : (Z, *) \longrightarrow (Z', *)$ in Ord_* satisfies $\phi(*) = *$ and $\phi(x) \leq \phi(y) \in Z'$ for any $x \leq y$ in Z. Let $\vec{\Gamma}_*$ denote the subcategory of Ord_* consisting of the objects $[n] = \{0 < 1 < 2 < ... < n\}$ (with basepoint 0) for any $n \geq 0$. Then, the category $\mathcal{C}$ determines a functor:

$$S(\mathcal{C}) : \vec{\Gamma}_* \longrightarrow SmCat_0 \qquad [n] \mapsto S_n\mathcal{C} \tag{2.2}$$

where $SmCat_0$ denotes the category of small categories with zero objects. The morphisms in $SmCat_0$ are functors that preserve zero objects. Given a morphism $\phi : [n] \longrightarrow [m]$ in $\vec{\Gamma}_*$, we have an induced functor:

$$S(\phi) : S_n\mathcal{C} \longrightarrow S_m\mathcal{C} \qquad (a_0, ...a_{n-1}) \mapsto (b_0, ..., b_{m-1}) \qquad b_j := \prod_{i \in \phi^{-1}(j), i<n} a_i \tag{2.3}$$

where in (2.3), for any $0 \leq j \leq m-1$, the cofibration $b_j = \prod_{i \in \phi^{-1}(j), i<n} a_i$ is the composition of the cofibrations a_i where i lies in the ordered set $\phi^{-1}(j)$ and $i < n$. In (2.3) it is understood that when $\phi^{-1}(j) \cap \{0, 1, 2, ..., n-1\}$ is empty, we set $b_j = 1$. It is clear that the functor $S(\mathcal{C}) : \vec{\Gamma}_* \longrightarrow SmCat_0$ in (2.2) extends to a functor from Ord_* to $SmCat_0$ that we continue to denote by $S(\mathcal{C}) : Ord_* \longrightarrow SmCat_0$. We are now ready to define the K-groups of $\mathcal{C}$ with respect to an ordered finite simplicial set Y with basepoint.

Definition 2.1. *Let $\mathcal{C}$ be a category as above and let $Y : \Delta^{op} \longrightarrow Ord_*$ be an ordered finite simplicial set with basepoint. Let $Sets_*$ denote the category of pointed sets. We consider the following composition of functors:*

$$S^Y(\mathcal{C}) : \Delta^{op} \xrightarrow{Y} Ord_* \xrightarrow{S(\mathcal{C})} SmCat_0 \xrightarrow{obj} Sets_* \tag{2.4}$$

where $obj : SmCat_0 \longrightarrow Sets_*$ is the functor that associates a category in $SmCat_0$ to its set of objects (with the zero object going to the basepoint). We consider the geometric realization $|S^Y(\mathcal{C})|$ of the pointed simplicial set $S^Y(\mathcal{C})$ in (2.4) and its loop space $\Omega|S^Y(\mathcal{C})|$. Then, we define K-theory groups $K_p^Y(\mathcal{C})$ of the category $\mathcal{C}$ of order Y to be the homotopy groups:

$$K_p^Y(\mathcal{C}) \coloneqq \pi_p(\Omega|S^Y(\mathcal{C})|) \qquad \forall\ p \geq 0 \tag{2.5}$$

We now show that homotopic maps of ordered simplicial sets determine identical morphisms on the K-groups defined above.

Proposition 2.2. *Let $\mathcal{C}$ be a small category with cofibrations. Let Y, $Y' : \Delta^{op} \longrightarrow Ord_*$ be ordered finite simplicial sets with basepoint. Let $f, g : Y \longrightarrow Y'$ be morphisms of simplicial objects of Ord_* that are simplicially homotopic. Then, f and g induce identical morphisms $K(f)_p = K(g)_p : K_p^Y(\mathcal{C}) \longrightarrow K_p^{Y'}(\mathcal{C})\ \forall\ p \geq 0$. In particular, if Y and Y' are simplicially homotopy equivalent as simplicial objects of Ord_*, $K_p^Y(\mathcal{C}) \cong K_p^{Y'}(\mathcal{C})$.*

Proof. Let $f_n, g_n : Y_n \longrightarrow Y'_n$ be the morphisms corresponding to f and g respectively at each level n. We are given that f and g are simplicially homotopic morphisms between simplicial objects of Ord_*. It follows that (see [4, § 8.3.11]) there are morphisms $h_{i,n} : Y_n \longrightarrow Y'_{n+1}$, $0 \leq i \leq n$, $n \geq 0$ in Ord_* such that $d^{Y'}_{0,n+1} h_{0,n} = f_n$ and $d^{Y'}_{n+1,n+1} h_{n,n} = g_n$ and

$$\begin{aligned} d^{Y'}_{i,n+1} h_{j,n} &= \begin{cases} h_{j-1,n-1} d^{Y}_{i,n} & \text{if } i < j \\ d^{Y'}_{i,n+1} h_{i-1,n} & \text{if } i = j \neq 0 \\ h_{j,n-1} d^{Y}_{i-1,n} & \text{if } i \geq j+1 \end{cases} \\ s^{Y'}_{i,n+1} h_{j,n} &= \begin{cases} h_{j+1,n+1} s^{Y}_{i,n} & \text{if } i \leq j \\ h_{j,n+1} s^{Y}_{i-1,n} & \text{if } i > j \end{cases} \end{aligned} \tag{2.6}$$

Here $d^Y_{i,n} : Y_n \longrightarrow Y_{n-1}$ (resp. $d^{Y'}_{i,n} : Y'_n \longrightarrow Y'_{n-1}$) and $s^Y_{i,n} : Y_n \longrightarrow Y_{n+1}$ (resp. $s^{Y'}_{i,n} : Y'_n \longrightarrow Y'_{n+1}$) for $0 \leq i \leq n$ are respectively the face and degeneracy maps of the simplicial object Y (resp. Y') of Ord_*. We now consider the simplicial sets $S^Y(\mathcal{C}) = \{S^Y(\mathcal{C})_n\}_{n\geq 0}$ and $S^{Y'}(\mathcal{C}) = \{S^{Y'}(\mathcal{C})_n\}_{n\geq 0}$ as defined in (2.4). By definition, for any $n \geq 0$, $S^Y(\mathcal{C})_n = obj(S(\mathcal{C})(Y_n))$ and $S^{Y'}(\mathcal{C})_n = obj(S(\mathcal{C})(Y'_n))$ along with induced maps $obj(S(\mathcal{C})(f_n)), obj(S(\mathcal{C})(g_n)) : S^Y(\mathcal{C})_n \longrightarrow S^{Y'}(\mathcal{C})_n$. Then, the induced maps $obj(S(\mathcal{C})(h_{i,n})) : S^Y(\mathcal{C})_n \longrightarrow S^{Y'}(\mathcal{C})_{n+1}$ define a simplicial homotopy between the two maps $S^f(\mathcal{C}) = \{obj(S(\mathcal{C})(f_n))\}_{n\geq 0}$ and $S^g(\mathcal{C}) = \{obj(S(\mathcal{C})(g_n))\}_{n\geq 0} : S^Y(\mathcal{C}) \longrightarrow S^{Y'}(\mathcal{C})$ of simplicial sets.

It now follows from the definitions in (2.5) that the induced morphisms $K(f)_p \coloneqq \pi_p(\Omega|S^f(\mathcal{C})|), K(g)_p \coloneqq \pi_p(\Omega|S^g(\mathcal{C})|) : K_p^Y(\mathcal{C}) = \pi_p(\Omega|S^Y(\mathcal{C})|) \longrightarrow \pi_p(\Omega|S^{Y'}(\mathcal{C})|) = K_p^{Y'}(\mathcal{C})$ on the homotopy groups are identical. □

Given small categories with cofibrations $\mathcal{C}$, $\mathcal{D}$, we consider the category $\mathcal{C} \times \mathcal{D}$. An object of $\mathcal{C} \times \mathcal{D}$ is a pair (C, D) where $C \in obj(\mathcal{C})$ and $D \in obj(\mathcal{D})$. For (C, D), $(C', D') \in obj(\mathcal{C} \times \mathcal{D})$, the collection of morphisms from (C, D) to (C', D') in $\mathcal{C} \times \mathcal{D}$ is given by $Hom_{\mathcal{C}}(C, C') \times Hom_{\mathcal{D}}(D, D')$. Given a small category with cofibrations $\mathcal{E}$, we now recall that a functor $F : \mathcal{C} \times \mathcal{D} \longrightarrow \mathcal{E}$ is said to be bi-exact if it satisfies the following two conditions (see, for instance, [2, Definition 4.2.1]):
(1) For any $C \in obj(\mathcal{C})$ (resp. $D \in obj(\mathcal{D})$), the functor $F(C, __) : \mathcal{D} \longrightarrow \mathcal{E}$ (resp. $F(__, D) : \mathcal{C} \longrightarrow \mathcal{E}$) is exact.
(2) Given cofibrations $C \rightarrowtail C'$ and $D \rightarrowtail D'$ in the categories $\mathcal{C}$ and $\mathcal{D}$ respectively, the canonical morphism from $F(C', D) \coprod_{F(C,D)} F(C, D')$ to $F(C', D')$ is a cofibration in $\mathcal{E}$.

Proposition 2.3. *Let $Y : \Delta^{op} \longrightarrow Ord_*$ be an ordered finite simplicial set with basepoint. Let $\mathcal{C}$, $\mathcal{D}$ and $\mathcal{E}$ be small categories with cofibrations and let $F : \mathcal{C} \times \mathcal{D} \longrightarrow \mathcal{E}$ be a bi-exact functor. Then, there exists a product structure:*

$$K_p^Y(\mathcal{C}) \times K_q^Y(\mathcal{D}) \longrightarrow K_{p+q}^Y(\mathcal{E}) \qquad \forall\ p, q \geq 0 \tag{2.7}$$

Proof. Given a bi-exact functor $F : \mathcal{C} \times \mathcal{D} \longrightarrow \mathcal{E}$, we consider the induced functors $F_n : S_n\mathcal{C} \times S_n\mathcal{D} \longrightarrow S_n\mathcal{E}$, $n \geq 0$ defined as follows:

$$\begin{array}{c} (0 = A_0 \twoheadrightarrow A_1 \twoheadrightarrow \cdots \twoheadrightarrow A_n) \times (0 = B_0 \twoheadrightarrow B_1 \twoheadrightarrow \cdots \twoheadrightarrow B_n) \in S_n\mathcal{C} \times S_n\mathcal{D} \\ \downarrow \\ (0 = F(A_0, B_0) \rightarrowtail F(A_1, B_1) \rightarrowtail \ldots \rightarrowtail F(A_n, B_n)) \in S_n\mathcal{E} \end{array} \tag{2.8}$$

We note that since F is bi-exact, the morphisms $F(A_k, B_k) \longrightarrow F(A_{k+1}, B_k)$ and $F(A_{k+1}, B_k) \longrightarrow F(A_{k+1}, B_{k+1})$ are cofibrations in $\mathcal{E}$ for each $k \geq 0$. Hence, each morphism $F(A_k, B_k) \longrightarrow F(A_{k+1}, B_{k+1})$ in (2.8) obtained by composing $F(A_k, B_k) \longrightarrow F(A_{k+1}, B_k)$ and $F(A_{k+1}, B_k) \longrightarrow F(A_{k+1}, B_{k+1})$ is a cofibration. Using the fact that F is bi-exact, we also see that $F(A_k, 0) = F(0, B_k) = 0$. Hence, it follows that (2.8) induces a morphism $obj(S_n\mathcal{C}) \wedge obj(S_n\mathcal{D}) \longrightarrow obj(S_n\mathcal{E})$ of pointed sets. Then, if we consider the ordered finite simplicial set $Y = \{Y_n\}_{n \geq 0}$, we have morphisms:

$$F_n^Y : obj(S(\mathcal{C})(Y_n)) \wedge obj(S(\mathcal{D})(Y_n)) \longrightarrow obj(S(\mathcal{E})(Y_n)) \qquad \forall\ n \geq 0 \tag{2.9}$$

From (2.9), it follows that we have a morphism $F^Y : S^Y(\mathcal{C}) \wedge S^Y(\mathcal{D}) \longrightarrow S^Y(\mathcal{E})$ of pointed simplicial sets. Passing to geometric realizations and taking loop spaces, we have an induced map $\Omega|S^Y(\mathcal{C})| \wedge \Omega|S^Y(\mathcal{D})| \longrightarrow \Omega|S^Y(\mathcal{E})|$. The result is now clear from the definitions in (2.5).

□

3 Hochschild homology and the Dennis trace map

We recall that a cyclic set is a contravariant functor from Connes' cyclic category ΔC to the category $Sets$ of sets (for details see, for instance, [1, § 6.1.2.1]). Given a small category $\mathcal{A}$, we can associate to it the cyclic set $CN(\mathcal{A}) = \{CN_n(\mathcal{A})\}_{n \geq 0}$ given by its cyclic nerve; in other words, for any $n \geq 0$, we set:

$$CN_n(\mathcal{A}) := \coprod_{(A_0, \ldots, A_n) \in obj(\mathcal{A})^{n+1}} Hom_{\mathcal{A}}(A_1, A_0) \times Hom_{\mathcal{A}}(A_2, A_1) \times \cdots \times Hom_{\mathcal{A}}(A_0, A_n) \tag{3.1}$$

By abuse of notation, given a small category $\mathcal{A}$, we will also let $CN(\mathcal{A}) = \{CN_n(\mathcal{A})\}_{n \geq 0}$ denote the underlying simplicial set of the cyclic set $CN(\mathcal{A})$.

Definition 3.1. *Let $\mathcal{C}$ be a small category with cofibrations and let $Y : \Delta^{op} \longrightarrow Ord_*$ be an ordered finite simplicial set with basepoint. Let k be a given field. We consider the composition of functors:*

$$CN(S^Y(\mathcal{C})) : \Delta^{op} \xrightarrow{\ Y\ } Ord_* \xrightarrow{\ S(\mathcal{C})\ } SmCat_0 \xrightarrow{\ CN\ } SSets \tag{3.2}$$

where $SSets$ denotes the category of simplicial sets. Let $|CN(S^Y(\mathcal{C}))|$ denote the geometric realization of the bisimplicial set $CN(S^Y(\mathcal{C}))$. Then, we define the Hochschild homologies $HH_p^Y(\mathcal{C})$ of the category $\mathcal{C}$ of order Y over the field k to be the homology groups:

$$HH_p^Y(\mathcal{C}) := H_{p+1}(|CN(S^Y(\mathcal{C}))|, k) \qquad \forall\ p \geq 0 \tag{3.3}$$

As noted before, the cyclic nerve of a small category is a cyclic set. As such, the bisimplicial set $CN(S^Y(\mathcal{C}))$ in (3.2) is actually a "cyclic $\times$ simplicial set" (i.e., a cyclic set in one coordinate and a simplicial set in the other; see, for instance, [2, Appendix A.6]). Taking the geometric realization first in the simplicial direction, we obtain a cyclic space whose geometric realization carries the structure of an S^1-space (see [1, § 7.1.4]). Then, we can consider the cyclic geometric realization $|CN(S^Y(\mathcal{C}))|^{cy}$ of $CN(S^Y(\mathcal{C}))$ which is given by the Borel space $|CN(S^Y(\mathcal{C}))|^{cy} \coloneqq ES^1 \times_{S^1} |CN(S^Y(\mathcal{C}))|$ (see [1, § 7.2.2]). Here ES^1 is any contractible space on which the topological group S^1 has a free action. We can define the cyclic homologies $HC_p^Y(\mathcal{C})$ of the category $\mathcal{C}$ of order Y to be the homology groups $HC_p^Y(\mathcal{C}) \coloneqq H_{p+1}(|CN(S^Y(\mathcal{C}))|^{cy}, k),\ \forall\ p \geq 0$.

Proposition 3.2. *Let $\mathcal{C}$ be a small category with cofibrations and let $Y : \Delta^{op} \longrightarrow Ord_*$ be an ordered finite simplicial set with basepoint. Then, the Hochschild and cyclic homologies of $\mathcal{C}$ of order Y fit into a long exact sequence:*

$$\ldots \longrightarrow HH_p^Y(\mathcal{C}) \longrightarrow HC_p^Y(\mathcal{C}) \longrightarrow HC_{p-2}^Y(\mathcal{C}) \longrightarrow HH_{p-1}^Y(\mathcal{C}) \longrightarrow \ldots \tag{3.4}$$

Proof. From [1, § 7.2.7], it follows that there exists a homotopy fibration $|CN(S^Y(\mathcal{C}))| \longrightarrow |CN(S^Y(\mathcal{C}))|^{cy} \longrightarrow BS^1$, where BS^1 is the classifying space of the topological group S^1. Hence, it follows from [1, § 7.2.10] that the long exact sequence given by the homology spectral sequence corresponding to this fibration gives us the long exact sequence in (3.4).

□

Proposition 3.3. *Let $\mathcal{C}$ be a small category with cofibrations. Let $Y,\ Y' : \Delta^{op} \longrightarrow Ord_*$ be ordered finite simplicial sets with basepoint. Let $f, g : Y \longrightarrow Y'$ be morphisms of simplicial objects of Ord_* that are simplicially homotopic. Then, f and g induce identical morphisms $H(f)_p = H(g)_p : HH_p^Y(\mathcal{C}) \longrightarrow HH_p^{Y'}(\mathcal{C}),\ \forall\ p \geq 0$. In particular, if $Y,\ Y'$ are simplicially homotopy equivalent as simplicial objects of Ord_*, $HH_p^Y(\mathcal{C}) \cong HH_p^{Y'}(\mathcal{C})$.*

Proof. For any fixed $n \geq 0$, we consider the simplicial set $CN_n(S^Y(\mathcal{C}))$ given by the composition:

$$CN_n(S^Y(\mathcal{C})) : \Delta^{op} \xrightarrow{Y} Ord_* \xrightarrow{S(\mathcal{C})} SmCat_0 \xrightarrow{CN_n} Sets \tag{3.5}$$

Then, as in the proof of Proposition 2.2, it follows that the maps of simplicial sets $CN_n(S^f(\mathcal{C}))$, $CN_n(S^g(\mathcal{C})) : CN_n(S^Y(\mathcal{C})) \longrightarrow CN_n(S^{Y'}(\mathcal{C}))$ induced by f and g respectively are simplicially homotopic.

If we consider the geometric realization of the cyclic $\times$ simplical set $CN(S^Y(\mathcal{C}))$ (resp. $CN(S^{Y'}(\mathcal{C}))$ in the simplicial direction, we obtain the cyclic space $\{|CN_n(S^Y(\mathcal{C}))|\}_{n\geq 0}$ (resp. $\{|CN_n(S^{Y'}(\mathcal{C}))|\}_{n\geq 0}$). Here, the space $|CN_n(S^Y(\mathcal{C}))|$ (resp. $|CN_n(S^{Y'}(\mathcal{C}))|$) is given by the geometric realization of the simplicial set $CN_n(S^Y(\mathcal{C})) = \{CN_n(S(\mathcal{C})(Y_m))\}_{m\geq 0}$ (resp. $CN_n(S^{Y'}(\mathcal{C})) = \{CN_n(S(\mathcal{C})(Y'_m)\}_{m\geq 0}$). From the above, it follows that the morphisms between the cyclic spaces $\{|CN_n(S^Y(\mathcal{C}))|\}_{n\geq 0}$ and $\{|CN_n(S^{Y'}(\mathcal{C}))|\}_{n\geq 0}$ induced respectively by f and g are homotopic in each degree. Hence, the morphisms between the geometric realizations of the cyclic $\times$ simplicial sets $CN(S^Y(\mathcal{C}))$ and $CN(S^{Y'}(\mathcal{C}))$ induced by f and g respectively are homotopic. It follows from (3.3) that the induced maps $H(f)_p, H(g)_p : HH_p^Y(\mathcal{C}) \longrightarrow HH_p^{Y'}(\mathcal{C})$ on the Hochschild homologies are identical.

□

From the proof of Proposition 3.3 and the definition $HC_p^Y(\mathcal{C}) \coloneqq H_{p+1}(|CN(S^Y(\mathcal{C}))|^{cy}, k)$, it is clear that given simplicially homotopic maps $f, g : Y \longrightarrow Y'$ as above, an analogous result holds for induced maps on cyclic homologies.

Proposition 3.4. *Let $Y : \Delta^{op} \longrightarrow Ord_*$ be an ordered finite simplicial set with basepoint. Let $\mathcal{C}$, $\mathcal{D}$ and $\mathcal{E}$ be small categories with cofibrations and let $F : \mathcal{C} \times \mathcal{D} \longrightarrow \mathcal{E}$ be a bi-exact functor. Then, there exists a product:*

$$HH_p^Y(\mathcal{C}) \otimes HH_q^Y(\mathcal{D}) \longrightarrow HH_{p+q+1}^Y(\mathcal{E}) \qquad \forall\ p, q \geq 0 \tag{3.6}$$

Proof. As in the proof of Proposition 2.3, for any $n \geq 0$, we have a functor $F_n : S_n\mathcal{C} \times S_n\mathcal{D} \longrightarrow S_n\mathcal{E}$ induced by $F : \mathcal{C} \times \mathcal{D} \longrightarrow \mathcal{E}$. Given $Y = \{Y_n\}_{n\geq 0} : \Delta^{op} \longrightarrow Ord_*$, it is clear that the functors F_n, $n \geq 0$ induce $F_n^Y : S(\mathcal{C})(Y_n) \times S(\mathcal{D})(Y_n) \longrightarrow S(\mathcal{E})(Y_n)$. Then, for any $m \geq 0$, we have a map:

$$\begin{array}{c} CN_m(F_n^Y) : CN_m(S(\mathcal{C})(Y_n)) \times CN_m(S(\mathcal{D})(Y_n)) \longrightarrow CN_m(S(\mathcal{E})(Y_n)) \\ (C_0 \to C_n \to C_{n-1} \to ... \to C_1 \to C_0) \times (D_0 \to D_n \to D_{n-1} \to ... \to D_1 \to D_0) \\ \Big\downarrow \\ (F(C_0, D_0) \to F(C_n, D_n) \to F(C_{n-1}, D_{n-1}) \to ... \to F(C_1, D_1) \to F(C_0, D_0)) \end{array} \tag{3.7}$$

The map of bisimplicial sets in (3.7) induces a morphism $|CN(F^Y)| : |CN(S^Y(\mathcal{C}))| \times |CN(S^Y(\mathcal{D}))| \longrightarrow |CN(S^Y(\mathcal{E}))|$ of geometric realizations. Finally, this gives us a product on homologies ($\forall\ p,\ q \geq 0$):

$$H_{p+1}(|CN(S^Y(\mathcal{C}))|, k) \otimes H_{q+1}(|CN(S^Y(\mathcal{D}))|, k) \longrightarrow H_{p+q+2}(|CN(S^Y(\mathcal{E}))|, k) \tag{3.8}$$

Comparing (3.8) with the definitions in (3.3), we obtain the map in (3.6).

□

Proposition 3.5. *Let $\mathcal{C}$ be a small category with cofibrations and let $Y : \Delta^{op} \longrightarrow Ord_*$ be an ordered finite simplicial set with basepoint. Then, for each $p \geq 0$, there is a morphism $D_p^Y : K_p^Y(\mathcal{C}) \longrightarrow HH_p^Y(\mathcal{C})$ from the K-groups of $\mathcal{C}$ of order Y to its Hochschild homology groups.*

Proof. Given a small category $\mathcal{A}$ in $SmCat_0$, for any $n \geq 0$, we have a map $CN_0(\mathcal{A}) \longrightarrow CN_n(\mathcal{A})$ of sets that takes any $A \in obj(\mathcal{A}) = CN_0(\mathcal{A})$ to $(A \xrightarrow{1} A \xrightarrow{1} A \xrightarrow{1} ... \xrightarrow{1} A) \in CN_n(\mathcal{A})$ (map $A \xrightarrow{1} A$ repeated n times). This gives us a morphism $CN_0(\mathcal{A}) \longrightarrow CN(\mathcal{A})$ of simplicial sets (where $CN_0(\mathcal{A})$ is treated as a constant simplicial set) and hence a morphism $CN_0 \longrightarrow CN$ of functors from $SmCat_0$ to $SSets$. Composing with $S(\mathcal{C}) \circ Y : \Delta^{op} \longrightarrow SmCat_0$, we have a morphism $D^Y : S^Y(\mathcal{C}) \longrightarrow CN(S^Y(\mathcal{C}))$ of functors from $\Delta^{op} \longrightarrow SSets$. The latter induces a map $|D^Y| : |S^Y(\mathcal{C})| \longrightarrow |CN(S^Y(\mathcal{C}))|$ of geometric realizations. Combining with the definitions in (2.5) and (3.3), we have a morphism $D_p^Y : K_p^Y(\mathcal{C}) \longrightarrow HH_p^Y(\mathcal{C})$ given by the composition:

$$K_p^Y(\mathcal{C}) = \pi_p(\Omega|S^Y(\mathcal{C})|) = \pi_{p+1}(|S^Y(\mathcal{C})|) \xrightarrow{\pi_{p+1}(|D^Y|)} \pi_{p+1}(|CN(S^Y(\mathcal{C}))|) \to H_{p+1}(|CN(S^Y(\mathcal{C}))|, k) = HH_p^Y(\mathcal{C})$$

□

References

[1] J.-L. Loday, Cyclic Homology, *Grundlehren der Mathematischen Wissenschaften*, **301**, Springer-Verlag, Berlin, 1998.

[2] R. McCarthy, The cyclic homology of an exact category. *J. Pure Appl. Algebra*, **93** (1994), no. 3, 251–296.

[3] F. Waldhausen, Algebraic K-theory of spaces. Algebraic and geometric topology (New Brunswick, N.J., 1983), 318–419, *Lecture Notes in Math.*, 1126, Springer, Berlin, 1985.

[4] C. Weibel, An introduction to homological algebra. *Cambridge Studies in Advanced Mathematics,* **38**. Cambridge University Press, Cambridge, 1994.